The complete chloroplast genome of *Limonium tetragonum* (Plumbaginaceae) isolated in Korea

Yongsung KIM, Hong XI¹ and Jongsun PARK^{1*}

Honam National Institute of Biological Resources, Mokpo 58762, Korea

¹InfoBoss Inc. and InfoBoss Research Center, Seoul 06278, Korea

(Received 27 June 2021; Revised 16 August 2021; Accepted 7 September 2021)

ABSTRACT: The chloroplast genome of *Limonium tetragonum* (Thunb.) Bullock, a halophytic species, was sequenced to understand genetic differences based on its geographical distribution. The cp genome of *L. tetragonum* was 154,689 bp long (GC ratio is 37.0%) and has four subregions: 84,572 bp of large single-copy (35.3%) and 12,813 bp of small single-copy (31.5%) regions were separated by 28,562 bp of inverted repeat (40.9%) regions. It contained 128 genes (83 protein-coding genes, eight rRNAs, and 37 tRNAs). Thirty-five single-nucleotide polymorphisms and 33 INDEL regions (88 bp in length) were identified. Maximum-likelihood and Bayesian inference phylogenetic trees showed that *L. tetragonum* formed a sister group with *L. aureum*, which is incongruent with certain previous studies, including a phylogenetic analysis.

Keywords: *Limonium tetragonum*, chloroplast genome, phylogenetic analysis, intraspecific variation, Plumbaginaceae

Limonium, belonging to Plumbaginaceae family, is a cosmopolitan halophytic genus, and a few species inhabit alkaline soil away from the coastal area (Kubitzki, 1993; Morgan and Funnell, 2018). *Limonium tetragonum* (Thunb.) Bullock, distributed in coastal areas of Korea and Japan, is a biennial halophytic species with radical leaves (8–15 × 1.5–3 cm) and yellow corolla (Owhi, 1965; Park, 2007; Lee et al., 2011; Park et al., 2020a). The crude extracts and solvent-partitioned fractions of whole plants of *L. tetragonum* display antioxidant (Lee et al., 2011) and anti-cancer activities (Kong et al., 2008); ethyl acetate soluble fraction of aerial parts exhibits anti-liver fibrosis (Kim et al., 2016); that of whole plants shows anti-alcohol toxicity (Kim et al., 2015); and its methanol extracts of whole plants present hepatoprotective activities (Yang et al., 2014). We sequenced *L. tetragonum* from the western coastal area of Korea for investigating intraspecific variations on chloroplast genomes with chloroplast of *L. tetragonum* isolated in eastern seashore of Korea.

Materials and Methods

Plant material

We collected *L. tetragonum* in Aphaedo island in Shinan-

gun, Jeollanam-do, Korea (34.8612N, 126.32629E). A voucher specimen and genomic DNA were deposited in the InfoBoss Cyber Herbarium (IN, the voucher number IB-00899).

DNA extraction and chloroplast genome determination

The total genomic DNA was extracted from fresh leaf by using a DNeasy Plant Mini Kit (QIAGEN, Hilden, Germany). Genome sequencing was performed using HiSeqX at Macrogen Inc., Korea, and *de novo* assembly was done by Velvet v1.2.10 (Zerbino and Birney, 2008) and GapCloser v1.12 (Zhao et al., 2011). Assembled sequences were confirmed by BWA v0.7.17 (Li, 2013) and SAMtools v1.9 (Li et al., 2009). All bioinformatic analyses were conducted in the Genome Information System (<http://geis.infoboss.co.kr/>) utilized in the previous studies (Kim et al., 2018, 2019a, 2019b, 2019c, 2021; Bum et al., 2020; Park et al., 2021c).

Genome annotation was conducted based on another *L. tetragonum* chloroplast (MW085088.1) with Geneious R11 v11.0.5 (Biomatters Ltd, Auckland, New Zealand). A circular map of *L. tetragonum* chloroplast genome was drawn using OGDRAW v1.31 (Greiner et al., 2019).

*Author for correspondence: starflr@infoboss.co.kr

Open Access

<http://e-kjpt.org>. © 2021 the Korean Society of Plant Taxonomists. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Identification of intraspecific variations

Single nucleotide polymorphisms (SNPs) and insertions and deletions (INDELs) were identified using the 'Find variations/SNPs' function implemented in the Geneious R11 v11.0.5 (Biomatters Ltd, Auckland, New Zealand) based on the pair-wise alignment of the two chloroplast genomes of *L. tetragonum* conducted by MAFFT 7.450 (Katoh and Standley, 2013). This method has been used in the previous studies (Kim et al., 2019a; Min et al., 2019a; Choi et al., 2021; Park et al., 2021b, 2021d). INDEL region was defined as the continuous INDELs.

Phylogenetic analysis

Maximum-likelihood (ML) and Bayesian inference (BI) phylogenetic trees were constructed based on the multiple sequence alignment of all available nine Plumbaginaceae chloroplast genomes and the seven chloroplast genomes of non-core Caryophyllales clade (Crawly and Hilu, 2012; Yao et al., 2019) by MAFFT v7.450 (Katoh and Standley, 2013). The chloroplast genome of *Nepenthes graciliflora* Elmer (1912) (Yao et al., 2019) was marked as outgroup species. The ML tree was reconstructed in IQ-TREE v1.6.12 (Nguyen et al., 2015) with 1,000 bootstrap repeats. In the ML analysis, a heuristic search was used with nearest-neighbor interchange branch swapping, GTR+F + R4 model determined as the best fit model by the ModelFinder implemented in IQ-TREE, and uniform rates among sites. All other options used the default settings. The posterior probability of each node was estimated by the BI using MrBayes v3.2.6 (Huelsenbeck and Ronquist 2001). The HKY85 model with gamma rates was used as a molecular model. A Markov chain Monte Carlo algorithm was employed for 1,100,000 generations, sampling trees every 200 generations, with four chains running simultaneously. Trees from the first 100,000 generations were discarded as burn-in.

Data availability

Chloroplast genome sequence can be accessed via accession number of MN044572 in GenBank of NCBI at <https://www.ncbi.nlm.nih.gov>. The associated BioProject, SRA, and Bio-Sample numbers are PRJNA737051, SAMN19678692, and SRR14793527, respectively.

Results and Discussion

The chloroplast genome of *L. tetragonum* (GenBank accession of MN044572) isolated in Korea is 154,689 bp long (GC ratio of 37.0%) and had four subregions: 84,572 bp of large single copy (35.3%) and 12,813 bp of small single copy (31.5%) regions are separated by 28,562 bp of inverted repeat (IR; 40.9%) (Table 1), which is similar to those of the previously reported *L. tetragonum* chloroplast genome (MW085088) (Table 1). It contained 128 genes (83 protein-coding genes [PCGs], eight rRNAs, and 37 tRNAs); 14 genes (five PCGs, four rRNAs, and five tRNAs) were duplicated in the IR regions, which is also same to those of the previously reported *L. tetragonum* chloroplast genome (MW085088) (Fig. 1, Table 1).

Based on the pair-wise sequence alignment with the previously reported *L. tetragonum* chloroplast genome (MW085088) (Darshetkar et al., 2021), 35 SNPs and 33 INDELs regions (88 bp in total) were identified. The longest INDEL region was 18-bp located between *trnT* and *trnL*. The inserted sequence was 18-bp repetitive sequences. In addition, two chloroplast genomes of *Limonium bicolor* (Bunge) Kuntze (Darshetkar et al., 2021) display extremely divergent manner, no SNP and 116 INDEL regions (15,840 bp in length), which is similar to the case of *Phedimus takesimensis* chloroplast genomes (129 SNPs and 112 INDEL regions (8,506 bp in length) (Park et al., in preparation).

Numbers of these intraspecific variations of *L. tetragonum* were smaller than those of *Pseudostellaria palibiniana* (Takeda) Ohwi (84 SNPs and 125-bp INDELs), *Pyrus ussuriensis* Maxim. (1,221 SNPs and 781-bp INDELs), *Goodyera schlechtendaliana* Rchb. f. (200 SNPs and 511-bp INDELs), *Gastrodia elata* Blume (324 SNPs and 630-bp INDELs) isolated in Korea; while they were larger than those of *Artemisia fukudo* Makino (seven SNPs and 12-bp INDELs) (Min et al., 2019b), *Fagus multinervis* Nakai (two SNPs and 2-bp INDELs) (Park and Oh, 2020), *Aconitum coreanum* (H. Lév.) Rapaics (five to 19 SNPs and 52-bp to 950bp INDELs) (Kim et al., 2019d), *Viburnum erosum* Thunb. (16 SNPs and 50-bp INDELs) (Choi et al., 2020), and *Veronica nakaiana* Ohwi (seven SNPs and 4-bp INDELs) (Lee et al., 2021). In

Table 1. List of two available chloroplast genomes of *Limonium tetragonum*.

GenBank accession	Length (bp)				GC contents				No. of genes		
	Whole	LSC	SSC	IR	Whole (%)	LSC (%)	SSC (%)	IR (%)	No. of PCGs	No. of tRNAs	No. of rRNAs
MN044572	154,689	84,572	13,013	28,562	37.0	35.3	31.5	40.9	83	37	8
MW085088	154,691	84,568	12,997	28,563	37.0	35.3	31.5	40.9	83	37	8

LSC, large single copy; SSC, small single copy; IR, inverted repeat ; PCG, protein-coding gene.

Fig. 1. Circular map of chloroplast genome of *Limonium tetragonum* isolated in Korea. Genes shown outside are transcribed clockwise, and those inside the circle are transcribed counter clockwise. Genes are color-coded to distinguish different functional groups. The dark grey and the light grey plot in the inner circle correspond to the GC content and AT content, respectively.

addition, seven PCGs contained at least one SNP or INDELs, among which two synonymous SNPs were found in *psbA* and *psaA* (Table 2) and four non-synonymous SNPs were identified in *rpl20*, *rpl32*, and *ycf2* located in the IR region (Table 2). The ratio of synonymous to non-synonymous SNPs was 0.5, which is different from the normal ratio, like *Chenopodium album* L. (Park et al., 2021a). One INDEL region was identified in the genic region of *rpoC2*, resulting the extension of *rpoC2* length in comparison to the previously sequenced chloroplast genome (MW085088) (Table 2).

Sixteen chloroplast genomes were used for constructing ML and BI phylogenetic trees. Both phylogenetic trees displayed that two *L. tetragonum*, collected from the East and West

coastal areas of Korean peninsula, respectively, were clustered with high bootstrap value (Fig. 2). The sister species of *L. tetragonum* is controversial: *L. tetragonum* formed a sister group with *Limonium aureum* (L.) Mill. in both trees (Fig. 2), whereas it was clustered with *L. bicolor* in the previous study using chloroplast genomes (Darshetkar et al., 2021). The close relative of *L. tetragonum* was not clearly confirmed because the phylogenetic study using chloroplast and nuclear loci could not determine the relationship with *Limonium sinense* (Girard) Kuntze, *Limonium tenellum* (Turczaninow) Kuntze, and *Limonium flexuosum* (Linnaeus) Kuntze (Koutroumpa et al., 2018) and the study with the complete chloroplast genome of *L. tetragonum* displayed that *L. bicolor* was sister species

Table 2. List of intraspecific variations identified from the two *Limonium tetragonum* chloroplast genomes.

No.	Type	Changed bases		Coordination		Type	Position	SNP type
		MN044572	MW085088	MN044572	MW085088			
1	SNP	G	A	485	485	Genic	psbA	Synonymous SNP
2	INDEL	-	AA	4,316	4,316-4,317	Intergenic	trnK-rps16	
3	SNP	G	A	4,549	4,551	Intergenic	trnK-rps16	
4	INDEL	T	-	6,256	6,258	Intergenic	rps16-trnQ	Frameshift
5	INDEL	-	AA	6,670	6,671-6,672	Intergenic	trnQ-psbK	
6	INDEL	-	AAA	7,214	7,217-7,219	Intergenic	psbK-psbI	
7	INDEL	-	T	7,866	7,872	Intergenic	trnS-trnG	Frameshift
8	INDEL	-	CGGGTCA	7,966	7,973-7,979	Intergenic	trnS-trnG	
9	INDEL	-	A	8,224	8,238	Intergenic	trnS-trnG	
10	INDEL	T	-	15,561	15,576	Intergenic	rps2-rpoC2	Frameshift
11	INDEL	-	T	15,898	15,912	Genic	rpoC2	
12	SNP	T	G	27,306	27,321	Intergenic	rpoB-trnC	
13	SNP	T	G	27,307	27,322	Intergenic	rpoB-trnC	Frameshift
14	SNP	C	G	27,308	27,323	Intergenic	rpoB-trnC	
15	SNP	C	A	27,309	27,324	Intergenic	rpoB-trnC	
16	SNP	C	A	27,310	27,325	Intergenic	rpoB-trnC	Frameshift
17	INDEL	-	A	28,271	28,286	Intergenic	trnC-petN	
18	INDEL	A	-	31,349	31,435	Intergenic	trnE-trnT	
19	INDEL	-	AA	31,472	31,487-31,488	Intergenic	trnE-trnT	Frameshift
20	SNP	C	A	31,768	31,785	Intergenic	trnE-trnT	
21	SNP	A	G	32,417	32,434	Intergenic	trnT-psbD	
22	INDEL	AAAG	-	36,846-36,849	36,863	Intergenic	psbZ-trnG	Synonymous SNP
23	SNP	G	A	42,109	42,122	Genic	psaA	
24	SNP	A	T	42,931	42,944	Intergenic	psaA-ycf3	
25	SNP	A	T	42,932	42,945	Intergenic	psaA-ycf3	Frameshift
26	INDEL	ATATATTATAT ATTATA	-	47,679-47,696	47,692	Intergenic	trnT-trnL	
27	INDEL	-	TAAT	47,873	47,868-47,871	Intergenic	trnT-trnL	
28	INDEL	ATTAAAA	-	48,687-48,693	48,686	Intronic	trnL	Frameshift
29	SNP	A	G	49,304	49,296	Intergenic	trnF-ndhJ	
30	INDEL	-	AAAA	55,296	55,288-55,291	Intergenic	atpB-rbcL	
31	INDEL	-	T	55,474	55,470	Intergenic	atpB-rbcL	Frameshift
32	INDEL	-	A	57,897	57,894	Intergenic	rbcL-accD	
33	INDEL	A	-	61,499	61,497	Intergenic	ycf4-cemA	
34	INDEL	-	TCTA	63,882	63,879-63,882	Intergenic	petA-psbJ	Frameshift
35	SNP	G	T	64,137	64,138	Intergenic	petA-psbJ	
36	SNP	A	T	64,138	64,139	Intergenic	petA-psbJ	
37	SNP	A	T	64,139	64,140	Intergenic	petA-psbJ	Frameshift
38	SNP	T	G	64,140	64,141	Intergenic	petA-psbJ	

Table 2. Continued

No.	Type	Changed bases		Coordination		Type	Position	SNP type
		MN044572	MW085088	MN044572	MW085088			
39	SNP	G	T	64,141	64,142	Intergenic	petA-psbJ	
40	SNP	A	T	64,142	64,143	Intergenic	petA-psbJ	
41	SNP	G	T	64,143	64,144	Intergenic	petA-psbJ	
42	SNP	A	C	64,146	64,147	Intergenic	petA-psbJ	
43	SNP	A	T	64,147	64,148	Intergenic	petA-psbJ	
44	SNP	A	C	64,148	64,149	Intergenic	petA-psbJ	
45	SNP	C	A	64,149	64,150	Intergenic	petA-psbJ	
46	SNP	A	T	64,150	64,151	Intergenic	petA-psbJ	
47	SNP	A	T	64,151	64,152	Intergenic	petA-psbJ	
48	SNP	A	C	64,152	64,153	Intergenic	petA-psbJ	
49	INDEL	T	-	65,511	65,512	Intergenic	psbE-petL	
50	INDEL	T	-	66,150	66,150	Intergenic	petL-petG	
51	INDEL	TA	-	68,210-68,211	68,209	Intergenic	rpl33-rps18	
52	INDEL	-	C	68,295	68,292	Intergenic	rpl33-rps18	
53	SNP	T	G	69,126	69,124	Genic	rpl20	Non-synonymous SNP
54	INDEL	CTTT	-	69,359-69,362	69,357	Intergenic	rpl20-rps12	
55	SNP	C	A	71,121	71,115	Intronic	clpP	
56	INDEL	A	-	75,382	75,376	Intergenic	psbH-petB	
57	INDEL	A	-	80,610	80,603	Intergenic	rps8-rpl14	
58	INDEL	-	AAAA	81,060	81,052-81,055	Intergenic	rpl14-rpl16	
59	SNP	C	A	86,943	86,939	Genic	ycf2	Non-synonymous SNP
60	INDEL	-	G	92,158	92,154	Intergenic	ycf2-trnL	
61	SNP	A	T	106,466	106,463	Intergenic	trnR-trnN	
62	SNP	A	T	115,749	115,746	Intergenic	ndhF-rpl32	
63	SNP	G	A	116,280	116,277	Genic	rpl32	Non-synonymous SNP
64	INDEL	-	T	116,808	116,805	Intergenic	rpl32-trnL	
65	INDEL	-	TTT	123,377	123,375-123,377	Intronic	ndhA	
66	SNP	T	A	132,796	132,797	Intergenic	trnN-trnR	
67	INDEL	-	C	147,096	147,097	Intergenic	trnL-ycf2	
68	SNP	G	T	152,319	152,321	Genic	ycf2	Non-synonymous SNP

SNP, single nucleotide polymorphism; INDEL, insertions and deletion.

(Darshetkar et al., 2021), however, number of taxa used in this study was limited due to lack of complete chloroplast genomes and *L. bicolor* was not included in the previous study. This incongruency of the phylogenetic relationship might be caused by different sequences in both studies. Moreover, the

supportive value of the clade of *L. tetragonum* and *L. bicolor* was 73%, which was lower than those (100%) in this study (Fig. 2), suggesting that *L. aureum* might be a sister species of *L. tetragonum*. Most of *Limonium* phylogenetic studies were conducted with the lack of East Asian *Limonium* species (Lledó

Fig. 2. Maximum-likelihood (ML) and Bayesian inference (BI) phylogenetic trees of 16 chloroplast genomes. Phylogenetic tree was drawn based on ML tree. The numbers above branches indicate support values of ML and BI trees, respectively.

et al., 2005; Malekmohammadi et al., 2017; Koutroumpa et al., 2018), requiring the additional phylogenetic studies including East Asian *Limonium* species.

ORCID: Yongsung KIM <http://orcid.org/0000-0002-5349-9226>;
Jongsun PARK <http://orcid.org/0000-0003-0786-4701>

Acknowledgments

This study was carried out with the support of the Honam National Institute of Biological Resources (HNIBR202101107).

Conflict of Interest

The authors declare that there are no conflicts of interest.

Literature Cited

- Bum, H. M., H. K Kim, N. S Kim, W. Kwon and J. Park. 2020. The complete chloroplast genome of *Douinia plicata* (Lindb.) Konstant. & Vilnet (Scapaniaceae, Jungermanniales). Mitochondrial DNA Part B Resources 5: 3680–3682.
- Choi, N. J., H. Xi and J. Park. 2021. A comparative analyses of the complete mitochondrial genomes of fungal endosymbionts in *Sogatella furcifera*, white-backed planthoppers. International Journal of Genomics 2021: 6652508.
- Choi, Y. G, N. Yun, J. Park, H. Xi, J. Min, Y. Kim and S.-H Oh. 2020. The second complete chloroplast genome sequence of the *Viburnum erosum* (Adoxaceae) showed a low level of intra-species variations. Mitochondrial DNA Part B Resources 5: 271–272.
- Crawley, S. S. and K. W. Hilu. 2012. Caryophyllales: Evaluating phylogenetic signal in *trnK* intron versus *matK*. Journal of Systematics and Evolution 50: 387–410.
- Darshetkar, A. M, S. Maurya, C. Lee, B. Bazarragchaa, G. Batdelger, A. Janchiv, E. J. Jeong, S. Choi, R. K Choudhary and S.-Y. Kim. 2021. Plastome analysis unveils Inverted Repeat (IR) expansion and positive selection in Sea Lavenders (*Limonium*, Plumbaginaceae, Limonioideae, Limonieae). PhytoKeys 175: 89–107.
- Greiner, S., P. Lehwark and R. Bock. 2019. OrganellarGenome-DRAW (OGDRAW) version 1.3. 1: Expanded toolkit for the graphical visualization of organellar genomes. Nucleic Acids Research 47: W59–W64.
- Huelsenbeck, J. P. and F. Ronquist. 2001. MRBAYES: Bayesian inference of phylogenetic trees. Bioinformatics 17: 754–755.
- Katoh, K. and D. M. Standley. 2013. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. Molecular Biology and Evolution 30: 772–780.
- Kim, M., H. Xi and J. Park. 2021. Genome-wide comparative analyses of GATA transcription factors among 19 *Arabidopsis* ecotype genomes: Intraspecific characteristics of GATA transcription factors. PLoS ONE 16: e0252181.
- Kim, N.-H., S. H. Sung, J.-D. Heo and E. J. Jeong. 2015. The extract of *Limonium tetragonum* protected liver against acute alcohol toxicity by enhancing ethanol metabolism and antioxidant enzyme activities. Natural Product Sciences 21: 54–58.
- Kim, N.-H., J.-D. Heo, T. B. Kim, J.-R. Rho, M. H. Yang and E. J.

- Jeong. 2016. Protective effects of ethyl acetate soluble fraction of *Limonium tetragonum* on diethylnitrosamine-induced liver fibrosis in rats. *Biological and Pharmaceutical Bulletin* 39: 1022–1028.
- Kim, Y., K.-I. Heo, S. Lee, and J. Park. 2018. Complete chloroplast genome sequence of the *Pseudostellaria longipedicellata* S. Lee, K. Heo & S.C. Kim (Caryophyllaceae). *Mitochondrial DNA Part B Resources* 3: 1296–1297.
- Kim, Y., Y. Chung and J. Park. 2019a. The complete chloroplast genome sequence of *Dysphania pumilio* (R. Br.) Mosyakin & Clemants (Amaranthaceae). *Mitochondrial DNA Part B Resources* 4: 403–404.
- Kim, Y., Y. J. Oh, K. Y. Han, G. H. Kim, J. Ko and J. Park. 2019b. The complete chloroplast genome sequence of *Hibiscus syriacus* L. ‘Mamonde’ (Malvaceae). *Mitochondrial DNA Part B Resources* 4: 558–559.
- Kim, Y., J. Park and Y. Chung. 2019c. Comparative analysis of chloroplast genome of *Dysphania ambrosioides* (L.) Mosyakin & Clemants understanding phylogenetic relationship in genus *Dysphania* R. Br. *Korean Journal of Plant Resources* 32: 644–668.
- Kim, Y., J.-S. Yi, J. Min, H. Xi, D. Y. Kim, J. Son, J. Park and J.-I. Jeon. 2019d. The complete chloroplast genome of *Aconitum coreanum* (H. Lévl.) Rapaics (Ranunculaceae). *Mitochondrial DNA Part B Resources* 4: 3404–3406.
- Kong, C.-S., Y. R. Um, J.-I. Lee, Y. A. Kim, J.-S. Lee and Y. Seo. 2008. Inhibition effects of extracts and its solvent fractions isolated from *Limonium tetragonum* on growth of human cancer cells. *Korean Journal of Biotechnology and Bioengineering* 23: 177–182.
- Koutroumpa, K., S. Theodoridis, B. H. Warren, A. Jiménez, F. Celep, M. Doğan, M. M. Romeiras, A. Santos-Guerra, J. M. Fernández-Palacios, J. Caujapé-Castells, M. Moura, M. M. de Sequeira and E. Conti. 2018. An expanded molecular phylogeny of Plumbaginaceae, with emphasis on *Limonium* (sea lavenders): Taxonomic implications and biogeographic considerations. *Ecology and Evolution* 8: 12397–12424.
- Kubitzki, K. 1993. Plumbaginaceae. In *The Families and Genera of Vascular Plants*. Vol. 2. Kubitzki, K., J. Rohwer and V. Bittrich (eds.), Springer-Verlag, Berlin. Pp. 523–530.
- Lee, J. I., C.-S. Kong, M. E. Jung, J. W. Hong, S. Y. Lim and Y. Seo. 2011. Antioxidant activity of the halophyte *Limonium tetragonum* and its major active components. *Biotechnology and Bioprocess Engineering* 16: 992.
- Lee, Y.-E., Y. Lee Y and S. Kim. 2021. A report of the second chloroplast genome sequence in *Veronica nakaiana* (Plantaginaceae), an endemic species in Korea. *Korean Journal of Plant Taxonomy* 51: 109–114.
- Li, H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv preprint at <https://arxiv.org/abs/1303.3997>.
- Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis and R. Durbin. 2009. The sequence alignment/map format and SAMtools. *Bioinformatics* 25: 2078–2079.
- Lledó, M. D., M. B. Crespo, M. F. Fay and M. W. Chase. 2005. Molecular phylogenetics of *Limonium* and related genera (Plumbaginaceae): Biogeographical and systematic implications. *American Journal of Botany* 92: 1189–1198.
- Malekmohammadi, M., H. Akhani and T. Borsch. 2017. Phylogenetic relationships of *Limonium* (Plumbaginaceae) inferred from multiple chloroplast and nuclear loci. *Taxon* 66: 1128–1146.
- Min, J., W. Kwon, H. Xi and J. Park. 2019a. The complete chloroplast genome of *Leucobryum juniperoides* (brid.) C. Müll. (Leucobryaceae, Bryophyta). *Mitochondrial DNA Part B Resources* 4: 2962–2963.
- Min, J., J. Park, Y. Kim and W. Kwon. 2019b. The complete chloroplast genome of *Artemisia fukudo* Makino (Asteraceae): Providing insight of intraspecies variations. *Mitochondrial DNA Part B Resources* 4: 1510–1512.
- Morgan, E., and K. Funnell. 2018. *Limonium*. In *Ornamental Crops*. Van Huylbroeck, J. (ed.), Springer, Cham. Pp. 513–527.
- Nguyen, L.-T., H. A. Schmidt, A. Von Haeseler and B. Q. Minh. 2015. IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* 32: 268–274.
- Ohwi, J. 1965. *Flora of Japan* (rev. ed.). Shibundo Co. Ltd., Tokyo, 344 pp.
- Park, C.-W.. 2007. *The Genera of Vascular Plants of Korea*. Flora of Korea Editorial Committee. Academy Publishing Co, Seoul, 1482 pp.
- Park, J. and S.-H. Oh. 2020. A second complete chloroplast genome sequence of *Fagus multinervis* Nakai (Fagaceae): Intraspecific variations on chloroplast genome. *Mitochondrial DNA Part B Resources* 5: 1868–1869.
- Park, J., J.-H. An, Y. Kim, D. Kim, B.-G. Yang and T. Kim. 2020a. Database of National Species List of Korea: The taxonomical systematics platform for managing scientific names of Korean native species. *Journal of Species Research* 9: 233–246.
- Park, J., J. Min, Y. Kim and Y. Chung. 2021a. The comparative analyses of six complete chloroplast genomes of morphologically diverse *Chenopodium album* L. (Amaranthaceae) collected in Korea. *International Journal of Genomics* 2021: 6643444.

- Park, J., S. Park, T. Jang, G. Kim and J.-H. Park. 2021b. The complete chloroplast genome of *Abeliophyllum distichum* f. *lilacinum* Nakai (Oleaceae) from the Chungbuk Province, Korea. Mitochondrial DNA Part B Resources 6: 1754–1756.
- Park, J., H. Xi, J. Son, H.T. Shin, H. Kang and S. Park. 2021c. The complete chloroplast genome of *Castanopsis sieboldii* (Makino) Hatus (Fagaceae). Mitochondrial DNA Part B Resources 6: 2743–2745.
- Park, J., H. Xi and Y. Kim. 2021d. The mitochondrial genome of *Arabidopsis thaliana* (Brassicaceae) isolated in Korea. Korean Journal of Plant Taxonomy. 51: 176–180.
- Yang, M. H., N.-H. Kim, J.-D. Heo, S. H. Sung and E. J. Jeong. 2014. Hepatoprotective effects of *Limonium tetragonum*, edible medicinal halophyte growing near seashores. Pharmacognosy Magazine 10: S563–S568.
- Yao, G., J.-J. Jin, H.-T. Li, J.-B. Yang, V. S. Mandala, M. Croley, R. Mostow, N. A. Douglas, M. W. Chase, M. J. M. Christenhusz, D. E. Soltis, P. S. Soltis, S. A. Smith, S. F. Brockington, M. J. Moore, T.-S. Yi and D.-Z. Li. 2019. Plastid phylogenomic insights into the evolution of Caryophyllales. Molecular Phylogenetics and Evolution 134: 74–86.
- Zerbino, D. R. and E. Birney. 2008. Velvet: Algorithms for de novo short read assembly using *de Bruijn* graphs. Genome Research 18: 821–829.
- Zhao, Q.-Y., Y. Wang Y, Y.-M. Kong, D. Luo, X. Li and P. Hao. 2011. Optimizing *de novo* transcriptome assembly from short-read RNA-Seq data: A comparative study. BMC Bioinformatics 12: S2.